


CELL REPRODUCTION

Cell Division

The process where one cell forms two identical daughter cells is called



Interphase is the first stage of the cell cycle.

- The cell matures, copies (replicates) its DNA, and prepares to divide.
- DNA stands for DeoxyriboNucleic Acid.
- **Replication** is a process when the cell makes a copy of its DNA in the nucleus.

The second stage of the cell cycle is called mitosis.

- During this stage the cell distributes all of the organelles and the DNA to each identical daughter cell.
- There are four phases during mitosis. They are: Prophase, Metaphase, Anaphase, and Telophase.

Components of DNA



To understand the phases of mitosis, you have to know the components of the DNA that is replicated in the process of mitosis. At the end of mitosis, one cell has become two identical cells, each with the same DNA. The stages of mitosis define the process by which the chromatin, the chromatic, the centromere, and the chromosome interrelate.

- The chromatin is the genetic material that has the instructions for how the cell is to be run. It is the actual DNA.
- ➤ The **chromatid** is the one half or one rod of the condensed chromatin.
- > The **chromosome** is the two daughter strands of a replicated chromatin that are condensed into a rod shape and connected by a centromere.
- ➤ The **centromere** is the point on the chromosome where

ids,





PREVIEW

Please <u>Sign In</u> or <u>Sign Up</u> to download the printable version of this worksheet












Sexual Reproduction and Meiosis

The majority of the species on Earth reproduce sexually.

The process that reduces the total number of chromosomes to half and forms the sex cells is called **meiosis**. The only cells that are created by meiosis are the sex cells. Sex cells are cells that make up the sperm (male) and the egg (female). There are two separate cell divisions during meiosis: meiosis 1 and meiosis 2. There are four phases in each cell division that have the same names as those from mitosis.

The end result of meiosis is four sex cells that have half of the genetic material of a mitotically produced cell.





The two sides of the double helix are made from a combination of molecules, one being a sugar and the other being a phosphate. They are called a nitrogen base when they are together.

Nitrogen Bases in DNA

There are four different nitrogen bases:

Adenine, Thymine, Guanine, and Cytosine. The nitrogen bases are referred to as A (Adenine), T (Thymine), G (Guanine), and C (Cytosine) when the topic is DNA.

A major point when talking about the nitrogen bases of DNA is that there is a specific pairing of the molecules. For example, A (Adenine) is always paired with T (Thymine) and G (Guanine) is always paired with C (Cytosine).



