

Motion



Name Class Date High velocity would describe a jet that is In order for motion to occur, there has to be a change in moving quickly in a certain direction. between two objects. The term most closely related to velocity is angle В A friction time c distance temperature **B** height **C** altitude D speed 3 When an object gains A car travels 100 miles in two hours at a speed, it is said to be speed of 50 miles per hour. Speed is accelerating. The term equal to the distance divided by 5 **PREVIEW** Please Sign In or Sign Up to download the printable version of this worksheet 7 Using the formula below, determine how True or false? many seconds it takes for it accelerate from 0 mph to 60 mph. final speed - initial speed B false acceleration time 2 seconds C 8 seconds 4 seconds D 15 seconds 9 Using the graph below, decide which sing the graph below, how fa statement is true. Distance vs. Time travel in the first 3 seconds? A the object is A 1 m **B** 3 m accelerating € 28 24 B the object is C 8 m decelerating **D** 15 m C the object is standing still D the object starts and stops



Motion



Name Class Date High velocity would describe a jet that is In order for motion to occur, there has to be a change in moving quickly in a certain direction. between two objects. The term most closely related to velocity is angle D В A friction time c distance temperature **B** height **C** altitude D speed 3 When an object gains A car travels 100 miles in two hours at a speed, it is said to be speed of 50 miles per hour. Speed is accelerating. The term equal to the distance divided by 5 **PREVIEW** D Please Sign In or Sign Up to download the printable version of this worksheet 7 Using the formula below, determine how True or false? many seconds it takes for it accelerate from 0 mph to 60 mph. B final speed - initial speed В false acceleration time 2 seconds C 8 seconds 4 seconds D 15 seconds Using the graph below, decide which sing the graph below, how fa statement is true. Distance vs. Time travel in the first 3 seconds? A the object is A 1 m **B** 3 m accelerating € 28 B the object is C 8 m decelerating **D** 15 m C the object is standing still D the object starts and stops